Seeing Olduvai Differently: Kikuletwa as a Modern Analog for Spring Refugia, Safe Sleeping Sites, and Hominin Landscape Use: a Commentary

IAN G. SMITH*
Departments Health, Environmental and Medical Sciences (HEMS), Faculty of Health, Environment and Medical Sciences, Bournemouth University, Fern Barrow, Poole, Dorset BH12 5BB, UNITED KINGDOM; ismith1@bournemouth.ac.uk

*corresponding author: Ian Smith; *ismith1@bournemouth.ac.uk*

submitted: 22 April 2026; revised 28 August 2026; accepted: 2 September 2026

Handling Editor in Chief: Karen Ruebens

ABSTRACT

Freshwater springs and associated wetlands are increasingly recognized as locally persistent resources within rift-basin paleolandscapes, and at Olduvai Gorge they have been linked to repeated concentrations of artifacts and fauna during dry phases. These reconstructions, however, often emphasize water and food availability, with less explicit attention to a major, underappreciated constraint, safe sleeping sites under nocturnal predation risk. This commentary uses Kikuletwa Hot Springs as a modern field prompt, alongside published reconstructions of spring and wetland use at Olduvai Bed I and Lowermost Bed II, to highlight the three-dimensional refuge structure that spring-oasis patches can support, especially woody cover and, where present, nearby rocky refuges generated by rift structure. By reframing springs as refugial nodes that bundle freshwater, edible plant resources, and refuge opportunities, this perspective helps separate activity locations from resting and sleeping needs within a wider pattern of movement and site use. It also leads to simple, testable predictions that can be applied to existing evidence and future work. The novelty is therefore not that springs attract hominins, but that spring-fed patches can be treated as bundled refuge structures, where freshwater, food opportunities, and nocturnal safety may be co-located, and where sleeping constraints may help organize daily ranging beyond the basin-floor record. Under this framing, archaeological evidence should occur more frequently near springs, wetlands, wooded patches, or other refuge features; these relationships should be weaker where deposits show greater water disturbance; and targeted survey of under-explored rocky ground and upland margins should reveal evidence that has so far been missed.

INTRODUCTION

Olduvai Gorge is frequently interpreted through the distribution of fossils and artifacts preserved in basin-floor and lake-margin sediments and landforms (Hay 1976; Leakey 1971). While these deposits are informative, they may not provide a complete picture of hominin landscape use. At Olduvai, hominin behavior is most visible archaeologically where tool use, butchery, or other artifact-rich activities left durable traces. Sleeping and resting behavior are far less likely to leave such clear signatures, even though they may have been equally important in structuring daily landscape use.

One recurring issue is that resource-rich lowlands can also be high-risk nocturnal environments for medium-sized terrestrial primates. For many primates, daily foraging on the ground is paired with a strong preference for elevated sleeping sites that reduce predation risk (Bidner et al. 2018; Hamilton III 1982; Lewis-Bevan et al. 2025). This safety constraint can shape how primates use landscapes, including where they choose to rest or sleep between foraging bouts, and it should be considered even if the Olduvai record is concentrated in lake-margin and basin-floor deposits.

Recent work at Olduvai has emphasized the importance of groundwater-fed springs and wetlands as locally persistent resources during climatic oscillations and has linked spring and wetland deposits to repeated concentrations of activity (Ashley et al. 2009; Ashley et al. 2010a; Ashley et al. 2010b; Cuthbert and Ashley 2014). The purpose of this short commentary is to add a behavioral layer to that framework, by treating springs not only as water points but as refugial nodes that can generate safe sleeping opportunities through associated tree cover. In short, the novelty here is not that springs attract hominins, but that spring-fed patches can form part of a three-dimensional refuge structure that constrains nocturnal safety and therefore shapes daily movement and site use.

Northern Tanzania contains multiple modern groundwater and wetland settings that can serve as partial ana-




Figure 1. Location of Kikuletwa Springs in relation to Mount Kilimanjaro and Mount Meru. The springs lie well away from the mountain flanks, illustrating the distal occurrence of groundwater discharge discussed in the text. Elevation contours, in meters above sea level, were generated by the author from the JAXA ALOS World 3D-30m (AW3D30) digital surface model, Version 3.2/3.1 (Japan Aerospace Exploration Agency Earth Observation Research Center 2021) (basemap: Esri World Imagery. Sources: Esri, DigitalGlobe, GeoEye, i-cubed, USDA FSA, USGS, AEX, Getmapping, Aerogrid, IGN, IGP, swisstopo, and the GIS User Community).

logs for aspects of Olduvai's rift-basin ecology (Ashley et al. 2010a). Kikuletwa is used here as the illustrative case because its spring discharge occurs well away from the mountain front, highlighting that reliable freshwater and associated tree cover can emerge distal to the recharge slopes, in locations that would otherwise appear exposed (Figure 1). This choice is therefore intended to clarify mechanism rather than to imply that Kikuletwa is the only or best analog in the region. A February 2026 visit to Olduvai Gorge and Kikuletwa provides the immediate field prompt for this perspective, but the argument is anchored in the published spring, wetland, and vegetation reconstructions for Bed I and Lowermost Bed II.

KIKULETWA AS A MODERN FIELD ANALOG

Kikuletwa Hot Springs, also known as Chemka, is a perennial spring oasis in northern Tanzania. The spring pools are bordered by dense riparian vegetation, including tall figs and palms, producing a compact patch of shade and elevated canopy within a broader, more open landscape (author observation). Hydrological field investigations at Chemka describe a high and relatively stable spring output throughout the year, including only modest fluctuations in water level and discharge, weak coupling to local precipitation, and a stable outlet temperature, consistent with a deep groundwater source and a sizable reservoir (Røhr et al. 2002). The spring discharge occurs distal to the Kilimanjaro recharge slopes, that is, well away from the mountain flanks, and the ground around the pools is relatively flat, which helps isolate canopy refuge as the primary three-dimensional structure at the site (Figures 2 and 3).

During a February 2026 visit, no baboons were observed, but vervet monkeys were observed resting high in the canopy above the spring pools and outflow. This qualitative observation is included for functional clarity rather than as a dataset, and it illustrates how groundwater discharge can generate a small, spatially concentrated refugial patch that combines reliable freshwater with elevated refuge structure in an otherwise open and exposed setting. Kikuletwa is now a recreational site with regular human activity, so present-day predator presence or absence cannot be regarded as representative of natural predator pressure.

Kikuletwa is therefore used here to visualize the behavioral logic of spring refugia. A spring-fed wetland can be more than a drinking point; it can also provide safe resting and sleeping options for ground-foraging primates. The next section summarizes the published evidence that comparable spring-fed wetland and wooded patches recur at Olduvai during Bed I and the earliest Bed II, where they have been linked to repeated hominin activity.



Figure 2. Riparian canopy at Kikuletwa Hot Springs (February 2026), illustrating the compact refuge structure associated with spring discharge.

SPRINGS AND WOODY PATCHES AT OLDUVAI

At Olduvai Gorge, multiple studies have reconstructed spring-fed freshwater oases and wetlands within a broader rift-basin mosaic that included saline or alkaline lake phases. In particular, spring and wetland deposits have been linked to repeated hominin activity during drier environmental phases, consistent with a freshwater spring refugia model (Ashley et al. 2009; Cuthbert and Ashley 2014). See also the mapped reconstruction of the wetland and spring sites in Ashley et al. (2009: Figure 5). In several reconstructions, springs are treated as especially important during drier phases, when shrinking lake-margin waters may have become increasingly saline or alkaline and less suitable for drinking.

Work at FLK North has explicitly described a freshwater oasis within savannah grassland settings, and spring-adjacent areas have been discussed as focal points for both hominin and carnivore behavior (Ashley et al. 2010a; Domínguez-Rodrigo et al. 2010). Freshwater spring settings were not necessarily low-risk environments—carnivore and crocodile activity is documented around FLK-Zinj and FLK NN3, reinforcing the importance of nearby protective vegetation or other refuge structure (Aramendi et al. 2017). Biomarker work at FLK Zinj, within a spring-fed wetland setting, indicates that freshwater, edible plant resources, and protective woody cover could be tightly bundled at meter scales, with butchery debris occurring within



Figure 3. Spring pool and outflow at Kikuletwa Hot Springs (February 2026), illustrating the oasis form discussed in the text.

or immediately adjacent to an isolated woodland patch near wetland vegetation (Magill et al. 2016), so a spring-fed patch can be a compact oasis that combines drinkable water, edible plants, and protective cover. By contrast, AGS (a Bed I site at Olduvai Gorge) shows that butchery and artifact concentrations could occur in a relatively open lakeside setting near freshwater, so the places where tools and bones accumulate are not always the places where hominins rested or slept (Stancampiano et al. 2022).

Most of this work explains spring-focused activity in terms of resources, drinkable water, and edible plants. The additional point emphasized here is that spring-associated woody patches can also provide refuge structure, including safer sleeping opportunities.

SAFE SLEEPING SITES AS A CONSTRAINT

In rift basins, groundwater discharge and local relief can share the same tectonic controls. Faulting can localize springs by creating conduits and barriers (Ashley et al. 2010b; Cuthbert and Ashley 2014), and it can also generate rocky scarps, outcrops, and steep ground (Hay 1976; Parisi et al. 2020). As a result, rift structure can produce a combined pattern of springs and nearby rocky refuges.

For baboons and other primates, sleeping site selection is strongly structured by predation avoidance, with cliffs, caves, and emergent trees typically preferred over open woodland trees when available (Hamilton III 1982). More recent work on leopard and baboon dynamics likewise emphasizes the role of sleeping sites in shaping risk and behavior (Bidner et al. 2018).

Applying this logic to Olduvai does not require claiming that hominins slept exclusively in trees. It suggests a spectrum of plausible refuges, including tall trees at spring oases, and, where available, fault-generated rocky ground and steeper slopes that can provide additional refuge options. Such relief need not occur at every activity locality. In an active rift setting, the availability of rocky refuges can also shift through time. The Olbalbal depression on the Ngorongoro flank is a reminder that faulting can rapidly reshape local topography, so rocky refuge options may have changed even where the present-day ground is relatively flat (Parisi et al. 2020).

This framing does not depend on any single explanation for how bones and artifacts accumulated at particular sites. Whatever the transport or deposition pathway, the main point is that interpretations of hominin habitat use should make room for a simple constraint. Where could vulnerable primates rest safely at night?

IMPLICATIONS FOR INTERPRETING OLDUVAI ASSEMBLAGES AND LAETOLI FOOTPRINTS

If spring-associated habitats are treated as refugial nodes, lake-margin artifact and faunal concentrations can be read as foraging and processing areas within a wider pattern of daily movement and site use, rather than as the only places where hominins spent most of their time. In this view, repeated use of spring and wetland locales is expected, but so is moving back and forth from safe refuges, including trees and nearby relief.

Paleobotanical syntheses indicate that spring-related wetlands and palm-rich woodlands were not unique to Olduvai but occurred as highly localized patches at Laetoli and Peninj as well, consistent with fault-controlled groundwater discharge and freshwater wet spots across northern Tanzania (Barboni 2014; Barboni et al. 2019). Work at Olduvai also shows that groundwater can be expressed as subtle seeps with diagnostic sedimentary signatures, suggesting that some spring-fed wet spots may be overlooked if attention is limited to the most obvious spring and wetland localities (Ashley et al. 2016). At a broader spatial scale, groundwater hydro-refugia modeling has mapped present-day "permanent" springs across the eastern branch of the East African Rift and indicates that many could remain productive through dry phases, reinforcing the likelihood that spring-fed nodes were widespread across the region (Cuthbert et al. 2017). This strengthens the inference that the wider region likely contained multiple spring-fed refugial patches beyond the best known localities, and that some potential nodes remain unrecognized because they have not been found, preserved, or exposed.

Bones and artifacts can be moved and re-concentrated after deposition, and similar looking patterns can be produced in different ways, so the refugial-node idea should be tested with spatial and sedimentary evidence rather than assumed from the positions of assemblages alone (Nel et al. 2021). At Olduvai, recent fluvial taphonomy work argues that many of the assemblages examined are largely in place, which means the spatial layout of bones and artifacts is more likely to be meaningful (Domínguez-Rodrigo et al. 2019).

The Laetoli trackways record a clear northerly movement direction. Miller et al. (2024) found that, while most footprint trails were oriented in random directions, all of the hominin trackways were directed northward, and suggested that this might reflect movement toward a water source. Published paleobotanical evidence indicates that localized spring-related wetlands and woodland occurred at Laetoli (Barboni 2014; Barboni et al. 2019). Modern hydrological work in the wider northern Tanzanian region also shows that groundwater discharge can emerge through spatially separated spring areas rather than at a single obvious source (Røhr et al. 2002). Within the framework proposed here, such localized groundwater-fed patches could have functioned as refugial nodes, combining water, vegetation, and refuge opportunities. An as yet unidentified spring-fed refugial node north of Laetoli therefore provides one plausible explanation for the observed northerly movement, although no such destination has yet been identified.

TESTABLE HYPOTHESES

This reframing leads to three clear, testable hypotheses that apply across rift-basin paleolandscapes.

1. If springs and refuge structure were important to hominins, archaeological evidence should occur more frequently near springs, wetlands, wooded patches,

or other safe refuge features.
2. Where archaeological deposits show greater water disturbance, their relationship to springs, wetlands, wooded patches, or other refuge features should be weaker.
3. If hominins also used refuge areas beyond the better-preserved basin floor, targeted survey of underexplored cliffs, rocky ground, and upland margins should reveal evidence that has so far been missed.

CONCLUSION

Kikuletwa is not presented as a direct analog for the geology or ecology of Bed I Olduvai. It is presented as a modern field prompt that makes the refugial structure of spring systems easier to appreciate. When combined with published reconstructions of spring and wetland use at Olduvai, it underpins a simple reframing—springs can be treated as refugial nodes that combine freshwater, plant resources, and elevated sleeping opportunities.

Making safe sleeping sites explicit as a constraint links primate ecology to rift-basin hydrology and taphonomy in a practical way. For expanded context on refugial landscape structure and behavioral constraints in an East African rift setting, see Smith and Reynolds (2025). It also provides simple predictions to check with maps, environmental evidence, and field survey. Above all, it is a reminder that basin-floor deposits may not tell the whole story of where early hominins rested, slept, and moved across the wider landscape.

ACKNOWLEDGMENTS

I thank Dr. Sally Reynolds for reading the draft, and for her helpful comments and encouragement, and I thank colleagues and guides in Tanzania for discussions during field visits to Olduvai Gorge and Kikuletwa Hot Springs in February 2026. Any errors of interpretation remain my own. No permits were required for the observations reported here, and no archaeological or paleontological materials were collected. No permissions were required for this commentary.

DATA AVAILABILITY

No new datasets were generated for this commentary. The argument is based on published sources cited in the text and on qualitative field observations. Photographs taken by the author during the February 2026 visit are available upon request.

AUTHOR CONTRIBUTION STATEMENT

IGS: conceptualization; methodology; software; validation; formal analysis; investigation; resources; data curation; writing – original draft; writing – review & editing; visualization; supervision; project administration; funding acquisition.

USE OF AI-ASSISTED TECHNOLOGIES

The author utilized ChatGPT (OpenAI) to improve readability and clarity during the writing process. After using ChatGPT, the author read and adjusted the text as needed and accepts full responsibility for the publication's content.



This work is distributed under the terms of a Creative Commons Attribution-NonCommercial 4.0 Unported License.

REFERENCES

Aramendi, J., Maté-González, M.A., Yravedra, J., Ortega, M.C., Arriaza, M.C., González-Aguilera, D., Baquedano, E., Domínguez-Rodrigo, M., 2017. Discerning carnivore agency through the three-dimensional study of tooth pits: revisiting crocodile feeding behaviour at FLK-Zinj and FLK NN3 (Olduvai Gorge, Tanzania). Palaeogeogr. Palaeoclimatol. Palaeoecol. 488, 93–102.

Ashley, G., Tactikos, J., Owen, R., 2009. Hominin use of springs and wetlands: paleoclimate and archaeological records from Olduvai Gorge (~1.79–1.74 Ma). Palaeogeogr. Palaeoclimatol. Palaeoecol. 272, 1–16.

Ashley, G.M., Barboni, D., Domínguez-Rodrigo, M., Bunn, H.T., Mabulla, A.Z., Diez-Martin, F., Barba, R., Baquedano, E., 2010a. Paleoenvironmental and paleoecological reconstruction of a freshwater oasis in savannah grassland at FLK North, Olduvai Gorge, Tanzania. Quat. Res. 74, 333–343.

Ashley, G.M., de Wet, C.B., Barboni, D., Magill, C.R., 2016. Subtle signatures of seeps: record of groundwater in a Dryland, DK, Olduvai Gorge, Tanzania. Depos. Rec. 2, 4–21.

Ashley, G.M., Domínguez-Rodrigo, M., Bunn, H.T., Mabulla, A.Z.P., Baquedano, E., 2010b. Sedimentary geology and human origins: a fresh look at Olduvai Gorge, Tanzania. J. Sed. Res. 80, 703–709.

Barboni, D., 2014. Vegetation of Northern Tanzania during the Plio-Pleistocene: a synthesis of the paleobotanical evidences from Laetoli, Olduvai, and Peninj hominin sites. Quat. Int. 322, 264–276.

Barboni, D., Ashley, G.M., Bourel, B., Arraiz, H., Mazur, J.-C., 2019. Springs, palm groves, and the record of early hominins in Africa. Rev. Palaeobot. Palynol. 266, 23–41.

Bidner, L.R., Matsumoto-Oda, A., Isbell, L.A., 2018. The role of sleeping sites in the predator-prey dynamics of leopards and olive baboons. Am. J. Primatol. 80, e22932.

Cuthbert, M.O., Ashley, G.M., 2014. A spring forward for hominin evolution in East Africa. PLoS One 9, e107358.

Cuthbert, M.O., Gleeson, T., Reynolds, S.C., Bennett, M.R., Newton, A.C., McCormack, C.J., Ashley, G.M., 2017. Modelling the role of groundwater hydro-refugia in East African hominin evolution and dispersal. Nat. Comm. 8, 15696.

Domínguez-Rodrigo, M., Baquedano, E., Barba, R., Uribelarrea, D., Gidna, A., 2019. The river that never was: fluvial taphonomy at Olduvai Bed I and II sites and its bearing on early human behavior. Quat. Int. 526, 26–38.

Domínguez-Rodrigo, M., Mabulla, A.Z.P., Bunn, H.T., Diez-Martin, F., Baquedano, E., Barboni, D., Barba, R., Domínguez-Solera, S., Sánchez, P., Ashley, G.M., Yravedra, J., 2010. Disentangling hominin and carni-

vore activities near a spring at FLK North (Olduvai Gorge, Tanzania). Quat. Res. 74, 363–375.

Hamilton III, W.J., 1982. Baboon sleeping site preferences and relationships to primate grouping patterns. Am. J. Primatol. 3, 41–53.

Hay, R.L., 1976. Geology of the Olduvai Gorge: A Study of Sedimentation in a Semiarid Basin. University of California Press, Berkeley.

Japan Aerospace Exploration Agency Earth Observation Research Center, 2021. ALOS Global Digital Surface Model (DSM): ALOS World 3D-30m (AW3D30), Version 3.2/3.1, 1.0 ed. Japan Aerospace Exploration Agency, Japan.

Leakey, M.D., 1971. Olduvai Gorge: Volume 3, Excavations in Beds I and II, 1960-1963. Cambridge University Press, Cambridge.

Lewis-Bevan, L., Hammond, P., Carvalho, S., Biro, D., 2025. Predictors of baboon sleep site selection in Gorongosa National Park. Am. J. Biol. Anthropol. 187, e70095.

Magill, C.R., Ashley, G.M., Domínguez-Rodrigo, M., Freeman, K.H., 2016. Dietary options and behavior suggested by plant biomarker evidence in an early human habitat. Proc. Nat. Acad. Sci. U.S.A. 113, 2874–2879.

Miller, C.K., McCann, R.A., Hatala, K.G., Musiba, C., DeSilva, J.M., 2024. Early hominin movement patterns at Laetoli, Northern Tanzania. PaleoAnthropology 2024:1, 139–147.

Nel, C., Bradfield, J., Lombard, M., Val, A., 2021. Taphonomic study of a modern baboon sleeping site at Misgrot, South Africa: implications for large-bodied primate taphonomy in karstic deposits. J. Paleolit. Archaeol. 4, 1–31.

Parisi, L., Stanistreet, I., Njau, J., Schick, K., Toth, N., Mai, P.M., 2020. Seismological investigations in the Olduvai Basin and Ngorongoro Volcanic Highlands (Western Flank of the North Tanzanian divergence). Seismol. Res. Lett. 91, 3286–3303.

Røhr, P.C., Ngana, J., Killingtveit, Å., 2002. Investigations of Chemka Spring, Kilimanjaro Region, Tanzania. In: Ngana, J.O. (Ed.), Water Resources Management: The Case of the Pangani River Basin: Issues and Approaches. Dar es Salaam University Press, Dar es Salaam, pp. 265–272.

Smith, I.G., Reynolds, S.C., 2025. Out of Afar: the first hominin migration? Long-term landscape changes in the Afar region and implications for hominin bipedalism [preprint]. Available at: https://doi.org/10.31223/X5TT85 [Accessed August 2026].

Stancampiano, L.M., Sistiaga, A., Del Val, D.U., Aramendi, J., Baquedano, E., Mabulla, A., Domínguez-Rodrigo, M., Magill, C.R., 2022. New site at Olduvai Gorge (AGS, Bed I, 1.84 Mya) widens the range of locations where hominins engaged in butchery. Sci. Rep. 12, 9794.